Please replace all prior claims in the application with the following:

1. (Currently Amended) A stabilized solid composition comprising containing a 4-amino-3-substituted-butanoic acid derivative and a stabilizer, the stabilizer being a humectant and effective against degradation of the 4-amino-3-substitutedbutanoic acid derivative due to lactam formation, the which comprises a 4-amino-3-substituted-butanoic acid derivative having the general formula

$$NH_2CH_2$$
 C CH_2COOH R_1 R_2

wherein,

R₁ is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R₂ is a monovalent group selected from:

- a straight or branched alkyl group of 3 8 carbon atoms;
- a straight or branched alkylene group of 3 8 carbon atoms;
- a straight or branched alkyl group of 3 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;
 - a cycloalkyl group of 3 8 carbon atoms;
- a cycloalkyl group of 3 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;
- a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

After Final Amendment/Required Submission for RCE—page 2 of 24



a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

PATENT PFIZER ANN ARBOR MI

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, and one or two of the unsubstituted methylene groups (-CH2-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an

After Final Amendment/Required Submission for RCE—page 3 of 24

alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

PATENT PFIZER ANN ARBOR MI

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, $-SO- or -S(O)_2-;$

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group; a condensed ring group formed by ortho-fusion of a phenyl ring

with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of

After Final Amendment/Required Submission for RCE—page 4 of 24

11 cont

T-111

5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

PATENT PFIZER ANN ARBOR MI

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a

After Final Amendment/Required Submission for RCE—page 5 of 24

Clark

5774-01-MR

Application No. 09/674,819

cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH2-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH2-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, a cycloalkanedienyl-methoxy group wherein one of the methylene groups (-CH2-) in said cycloalkanedienyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, or a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring wherein said cycloalkanedienyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

After Final Amendment/Required Submission for RCE—page 6 of 24

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or **-SS-**;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

PATENT PFIZER ANN ARBOR MI

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

O1,

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

After Final Amendment/Required Submission for RCE—page 7 of 24

of cont

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

PATENT PFIZER ANN ARBOR MI

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkyl group, an alkyl group, an alkyl group, an analyl group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a

After Final Amendment/Required Submission for RCE—page 8 of 24

C' cont

hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group; a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group; and

a-humectant for manufacturing a pharmaceutical preparation.

- 2. (Currently Amended) The stabilized solid composition containing a 4 amino 3 substituted butanoic acid derivative as claimed in of claim 1 wherein said humectant comprises one or more of the compounds selected from ethylene glycol, propylene glycol, butylene glycol, sorbitol, and glycerol and an a lower aliphatic acid ester thereof of glycerol.
- 3. (Currently Amended) The stabilized solid composition containing a 4-amino-3 substituted butanoic acid derivative as claimed in of claim 1 wherein said humectant is ethylene glycol.

After Final Amendment/Required Submission for RCE—page 9 of 24

y cont

4. (Currently Amended) The stabilized solid composition containing a 4-amine 3 substituted butanoic acid derivative as claimed in of claim 1 wherein said humectant is propylene glycol.

PATENT PFIZER ANN ARBOR MI

- 5. (Currently Amended) The stabilized solid composition containing a 4-amine 3-substituted butaneic acid derivative as claimed in of claim 1 wherein said humectant is butylene glycol.
- 6. (Currently Amended) The stabilized solid composition containing a 4-amino-3-substituted butanoic acid derivative as claimed in of claim 1 wherein said humectant is glycerol or an a lower aliphatic acid ester thereof of glycerol.
- 7. (Currently Amended) The stabilized solid composition containing a 4-amino-3-substituted butanoic acid derivative as claimed in of claim 1 wherein said humectant is sorbitol.
- 8. (Currently Amended) The stabilized solid composition containing a 4-amino-3-substituted-butanoic acid derivative as claimed in of claim 1 wherein a total amount of said humectant is 0.01 - 25% by weight relative to the 4-amino-3-substitutedbutanoic acid derivative.
- 9. (Currently Amended) The stabilized solid composition containing a 4-amino 3-substituted butanoic acid derivative as claimed in of claim 18 20 wherein a total amount of said humectant is 0.01 - 25% by weight relative to a total amount of the 4-amino-3-substituted-butanoic acid derivative and an the auxiliary agent for manufacturing a pharmacoutical proparation.
- 10. (Currently Amended) The stabilized solid composition containing a 4-amino 3-substituted butanoio acid derivative as claimed in any-of claims 1 to 9 wherein it is a solid pharmaceutical preparation of the 4-amino-3-substituted-butanoic acid

After Final Amendment/Required Submission for RCE—page 10 of 24

ent

5774-01-MR

Application No. 09/674,819

derivative is gabapentin, pregabalin, baclofen, 3-aminomethyl-4-cyclohexyl-butanoic acid, 3-aminomethyl-5-cyclohexyl pentanoic acid, 3-aminomethyl-4-phenyl-butanoic acid or 3-aminomethyl-5-phenyl-pentanoic acid and the solid composition is a pharmaceutical preparation.

PATENT PFIZER ANN ARBOR MI

- 11. (Currently Amended) The stabilized solid composition containing a 4 amino-3-substituted-butanoic acid-derivative as claimed in of claim 10 wherein it is a solid the pharmaceutical preparation in the is a dosage form of tablets, powders, granules or capsules.
- 12. (Currently Amended) A process for the preparation of preparing a solid composition containing a 4-amino-3-substituted-butanoic acid derivative having the general formula

wherein.

R₁ is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R₂ is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3-8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an

After Final Amendment/Required Submission for RCE—page 11 of 24

alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

PATENT PFIZER ANN ARBOR MI

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms:

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-;

After Final Amendment/Required Submission for RCE—page 12 of 24

1 ent

PATENT PFIZER ANN ARBOR MI

5774-01-MR Application No. 09/674,819

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, and one or two of the unsubstituted methylene groups (-CH2-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, $-SO- or -S(O)_2-;$

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of

After Final Amendment/Required Submission for RCE—page 13 of 24

5 - 8 carbon atoms, one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

PATENT PFIZER ANN ARBOR MI

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH2-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)2-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH2-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)2- and one or two of the unsubstituted methylene groups (-CH2-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is

After Final Amendment/Required Submission for RCE—page 14 of 24

PATENT PFIZER ANN ARBOR MI

5774-01-MR Application No. 09/674,819

mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH2-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH2-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, a cycloalkanedienyl-methoxy group wherein one of the methylene groups (-CH2-) in said cycloalkanedienyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, or a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring wherein said cycloalkanedienyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a

C'ent

After Final Amendment/Required Submission for RCE—page 15 of 24

carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

PATENT PFIZER ANN ARBOR MI

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl

After Final Amendment/Required Submission for RCE—page 16 of 24

group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

PATENT PFIZER ANN ARBOR MI

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2- group and one or more of the unsubstituted methylene groups (-CH2-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted

After Final Amendment/Required Submission for RCE—page 17 of 24

methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

wherein said process comprises comprising combining the 4-amino-3-substituted-butanoic acid derivative with a stabilizer, the stabilizer being a humectant for manufacturing a pharmaceutical preparation and effective against degradation of the 4-amino-3-substituted-butanoic acid derivative due to lactam formation.

13. (Currently Amended) The process as claimed in of claim 12 wherein said humectant comprises one or more of the compounds selected from ethylene glycol, propylene glycol, butylene glycol, sorbitol, and glycerol and an a lower aliphatic acid ester thereof of glycerol.

After Final Amendment/Required Submission for RCE—page 18 of 24

C) cont

14. (Currently Amended) The process as claimed in any-of claims 12 to 13 wherein said solid composition is a solid pharmaceutical preparation of gabapentin, pregabalin, baclofen, 3-aminomethyl-4-cyclohexyl-butanoic acid, 3-aminomethyl-5-cyclohexyl pentanoic acid, 3-aminomethyl-4-phenyl-butanoic acid or 3-aminomethyl-5-phenyl-pentanoic acid.

PATENT PFIZER ANN ARBOR MI

- 15. (Currently Amended) The process as claimed in of claim 14 wherein a solid the pharmaceutical preparation containing a 4 amino-3 substituted butanoic acid derivative is a pharmaceutical preparation in the is a dosage form of tablets, powders, granules or capsules.
- 16. (Currently Amended) A stabilized The solid composition containing a 4 amino 3 substituted butanoic-acid derivative as claimed in of claim 1 wherein it is further combined with further comprising a neutral amino acid.
- 17. (Currently Amended) The stabilized solid composition containing a 4 amino 3 substituted butanoic acid derivative as claimed in of claim 16 wherein said neutral amino acid is one or more of the neutral amino acids selected from L-leucine, L-isoleucine, L-valine, L-alanine, D-leucine, D-isoleucine, D-valine, D-alanine, DL-leucine, DL-isoleucine, DL-isoleucine, DL-valine, DL-alanine and glycine.
 - 18. (Canceled)
 - 19. (Canceled)
- 20. (Currently Amended) A stabilized The solid composition according to of Claim 1, wherein the 4 amino 3 substituted butanoic acid derivative is further combined with comprising an auxiliary agent.
- 21. (Currently Amended) The process as elaimed in of Claim 12, further comprising adding an auxiliary auxiliary agent to the solid composition containing the 3-substituted butanoic acid derivative.

After Final Amendment/Required Submission for RCE—page 19 of 24

C'ent

(New) A solid composition comprising gabapentin and a stabilizer, the 22. stabilizer being a humectant and effective against degradation of gabapentin due to lactam formation.

PATENT PFIZER ANN ARBOR MI

- (New) The solid composition of claim 22, wherein said stabilizer is one or 23. more of ethylene glycol, propylene glycol, butylene glycol, sorbitol, glycerol, or a lower aliphatic acid ester of glycerol.
- (New) The solid composition of claim 22, wherein said stabilizer is 24. propylene glycol.
- (New) A process for preparing a solid composition comprising combining 25. gabapentin with a stabilizer, the stabilizer being a humectant and effective against degradation of gabapentin due to lactam formation.
- (New) The process of claim 25, further comprising combining gabapentin 26. with one or more of ethylene glycol, propylene glycol, butylene glycol, sorbitol, glycerol, or a lower aliphatic acid ester of glycerol.
- (New) The process of claim 25, further comprising combining gabapentin 27. with propylene glycol.